

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C11D 3/39, 3/395	A2	(11) International Publication Number: WO 98/03621 (43) International Publication Date: 29 January 1998 (29.01.98)
(21) International Application Number: PCT/US97/12215 (22) International Filing Date: 18 July 1997 (18.07.97) (30) Priority Data: 60/022,334 24 July 1996 (24.07.96) US (71) Applicant (for all designated States except US): THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): OFOSU-ASANTE, Kofi [GH/US]; 1535 Cedar Avenue, Cincinnati, OH 45224 (US). HUTTON, Howard, David [US/US]; 2745 Jessica Court, Loveland, OH 45140 (US). (74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: METHOD FOR STAIN REMOVAL ON FABRIC WITH DETERGENT COMPOSITIONS CONTAINING BLEACH (57) Abstract The present invention is in the field of liquid or gel detergents. More specifically, the invention relates to a method for removing stains and/or odors and for disinfecting fabrics by applying a liquid or gel detergent composition to the fabric and subjecting the fabric to microwaves. This method provides enhanced cleaning and improved stain removal. These cleaning compositions comprise a bleaching agent, preferably diacyl peroxide in a liquid or gel detergent formulation. These compositions in combination with the method of microwaving are also very effective in disinfecting and deodorizing fabrics.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

METHOD FOR STAIN REMOVAL ON FABRIC WITH DETERGENT COMPOSITIONS CONTAINING BLEACH

TECHNICAL FIELD

The present invention is in the field of bleaching compositions, preferably liquid or gel detergents. More specifically, the invention relates to a method for removing stains, odors and for disinfecting fabric by applying a bleaching composition, preferably a liquid or gel detergent composition, and subjecting the fabric to microwaves. This method provides enhanced cleaning and improved stain removal. These cleaning compositions comprise a bleaching agent, preferably diacyl peroxide in a liquid or gel detergent formulation.

BACKGROUND OF THE INVENTION

Detergents used for washing fabric in the home or institution have long been known. The body of art pertaining to fabric cleaning is immense and encompasses many formulations designed for stain removal, many including bleaches.

However, consumers continue to experience problems with stain removal on fabric. In particular, formulators have experienced difficulties in formulating detergents which remove both hydrophobic and hydrophilic stains. Typically for stain removal, formulators have turned to chlorine bleach or sources of hydrogen peroxide and bleach activators.

Numerous substances have been disclosed in the art as effective bleach activators. One widely-used bleach activator is tetraacetyl ethylene diamine (TAED). TAED provides effective hydrophilic cleaning especially on beverage stains, but has limited performance on hydrophobic stains, dingy stains and body soils. Another type of activator, such as nonanoyloxybenzenesulfonate (NOBS) and other activators which generally comprise long chain alkyl moieties, is hydrophobic in nature and provides excellent performance on dingy stains and carotenoid stains.

It would seem that a combination of bleach activators, such as TAED and NOBS, would provide an effective detergent composition which would perform well on both hydrophilic and hydrophobic soils and stains. However, many of the hydrophilic activators developed thus far, including TAED, have been found to have limited efficacy, especially at laundry liquor temperatures below 60°C. Another consideration in the development of consumer products effective on both types of soils is the additional costs associated with the inclusion of two or more bleach activators. Accordingly, it is of

substantial interest to the manufacturers of bleaching systems to find a less expensive type of bleaching activator.

Chlorine bleaches are effective for stain and/or soil removal. While chlorine bleach is a very effective cleaning agent, it is not compatible with a variety of detergent ingredients and may require lengthy soaking time in which the bleach and the stained fabric must remain in contact to ensure stain removal.

Another known bleaching source is diacyl peroxides (DAPs). Although DAPs have been disclosed for use in the laundry and anti-acne area, they have had limited success in liquid or automatic dishwashing detergent area. In the laundry field certain diacyl peroxides have been disclosed as beneficial in cleaning tea stains from fibrous material.

Another problem facing formulators is stability of the bleaching agents and other individual ingredients over time, especially in liquid products. This is particularly true for diacyl peroxides in alkaline conditions.

As a consequence to the above-identified problems, there has been a substantial amount of research to develop bleaching systems which are stable and effective in liquid formulations and in which the amount of soaking time needed to remove stains is greatly reduced.

By the present invention, it has now been discovered that certain bleaching agents are stable and perform very well on stains when the bleaching agent is applied to a fabric and subjected to microwaves. Accordingly, the present invention solves the long-standing need for an inexpensive bleaching system which performs efficiently and effectively under mixed soil load conditions, especially mixtures of hydrophobic and hydrophilic soils.

The detergent compositions provided herein have the property of removing stains, especially tea, fruit juice and carotenoid stains objected to by the consumer from fabrics when exposed to microwaving from any typical household or commercial microwave. The compositions have other cleaning benefits in addition to stain removal advantages such as deodorizing and disinfecting. Thus, provided herein is a method of removing stains from a fabric by applying a bleaching compositions and exposing the stained fabric to microwaves to provide powerful cleaning of wide-ranging stains while retaining the advantages of a stable, mild product matrix.

SUMMARY OF THE INVENTION

The invention herein is a method for treating fabrics comprising the steps of:

- a) contacting, in the presence of water or a solvent which generates heat under microwave radiation, a fabric with a treating composition comprising an effective bleaching agent;
- b) subjecting the fabric to microwaves for a sufficient period to effectively treat the fabric.

There are also provided embodiments wherein additional components, especially surfactant, solvent, clay, polycarboxylate thickeners, baking soda, carbonates, phosphates, hydrobenzoic acid, dicarboxylic acid, siloxanes, perfumes, bleach catalysts, water, and mixtures thereof. Said compositions have a pH of from about 3 to about 13.

Examples of bleaching agents which are highly preferred for use herein included: diacyl peroxide, a source of hydrogen peroxide with or without a bleach activator, chlorine bleach and mixtures thereof.

Although this method can be employed to assist in bleaching the fabric for any desired purpose, examples of likely uses include one or more of the following: stain removal, stain reducing, deodorizing, or disinfecting. The fabric after being treated with the bleach composition is then subjected to microwaves for a sufficient time to "treat" the fabric to the desired end result, i.e. stain reduction, deodorizing, disinfecting. Typically a "sufficient time" in step b) is from about 1 second to about 2 minutes, preferably from about 10 seconds to about 45 seconds.

The method herein when employed to remove or reduce stains is highly effective on hydrophobic stains such as tomato stains and dingy stains. However, the method can also be employed with very good results on hydrophilic stains such as tea stains. Other example wherein the method is highly effective include the reduction or removal of hydrophobic stains on hydrophobic fabrics and the reduction or removal of hydrophilic stains on hydrophilic fabrics. An example of hydrophobic stains on hydrophobic substrates include tomato sauce on acrylic. An example of hydrophilic stains on hydrophilic substrates include tea stains on cotton. Included below are examples of treating composition which can be employed in the method of this invention.

- 1) A sprayable liquid or sprayable gel detergent compositions especially effective at cleaning stains, deodorizing, and/or disinfecting fabric, comprising by weight:
 - (a) from about 0.1% to about 60%, preferably 0.1 to about 40%, more preferably from about 0.3% to about 10%, of a bleaching agent selected from the group consisting of:
 - i) diacyl peroxide having the general formula:



wherein R and R1 can be the same or different and are hydrocarbyls, preferably no more than one is a hydrocarbyl chain of longer than ten carbon atoms, more preferably at least one has an aromatic nucleus;

- ii) a source of hydrogen peroxide;
 - iii) a source of hydrogen peroxide and a bleach activator;
 - iv) a chlorine bleach, preferably hypochlorite; and
 - v) mixtures thereof;
- (b) from 0% to about 95% of a non-aqueous solvent;
- (c) from 0% to about 50%, preferably about 2% to about 45%, more preferably from about 6% to about 30%, still more preferably from about 6% to about 12%, of a surfactant; and
- (d) from 0% to about 7%, preferably from about 0.5% to about 5%, more preferably from about 0.75% to about 3%, of a thickener.

2) A gel detergent composition especially effective at cleaning stains and tough food soil from fabrics, comprising by weight:

- (a) from about 0.1% to about 10%, preferably 0.1 to about 8%, more preferably from about 0.3% to about 5%, of a diacyl peroxide having the general formula:



wherein R and R1 can be the same or different and are hydrocarbyls, preferably no more than one is a hydrocarbyl chain of longer than ten carbon atoms, more preferably at least one has an aromatic nucleus;

- (b) from about 2% to about 45%, preferably from about 6% to about 40%, preferably from about 6% to about 12%, of a surfactant; and
- (c) from 0.5% to about 5%, preferably from about 0.75% to about 3%, of a thickener, preferably a clay or polycarboxylate thickener, more preferably Polygel® thickener;

said composition having a neat pH of from about 3 to about 10, preferably from about 6 to about 9; and such that said diacyl peroxide remains undissolved in said compositions and acts as an abrasive. The compositions herein are also effective at deodorizing and disinfecting fabrics.

For compositions which provide both stain removal and tough food cleaning as exemplified by 2) above, solvents are preferably not present. Without being limited by theory, it is believed that the presence of solvents may act to dissolve the diacyl peroxide and thus reduce the abrasive character of the composition. Thus, for an abrasive, tough

food cleaning formulation, it is preferable that the solvent be limited to no more than about 4%, more preferably no more than about 2%, by weight of the composition.

For sprayable compositions as exemplified by 1) above, solvent may be present with any of the above identified bleaching agents. However, when diacyl peroxide is present as the bleaching agent, it is preferable to have from about 2% to about 80% of a solvent capable of dissolving the diacyl peroxide. Suitable solvents are preferably selected from the group consisting of N-alkyl pyrrolidones, such as N-ethyl pyrrolidone, diacetone alcohol, alkyl ethers, cyclic alkyl ketones, and mixtures thereof. Amines, ethers and low molecular weight primary and secondary alcohols (about C₁-C₆) are preferably not present. Without being limited by theory, it is believed that the presence of these compounds may introduce stability problems. Thus, when diacyl peroxide and solvent are present in the compositions of this invention, it is further preferable that the amount of amine, ether, or primary or secondary alcohol be limited to no more than about 5%, preferably no more than about 3%, by weight of the composition.

Before microwaving, preferably water should also be present. The substrate may be wetted or dampened by water before or after application of the bleaching composition. Preferably, the bleaching composition comprises water. The compositions herein may, therefore, additionally comprise from about 0.1% to about 99.5%, more preferably from about 60-95%, even more preferably from about 80% to about 95%, by weight of the composition of water. Alternatively, a suitable solvent can be used which acts as a substitute for the water. Without being limited by theory, it is believed that the water or solvent readily generates heat when subjected to microwave radiation, which in turn, activates the bleaching agent. Suitable solvents, like water, are those with high dielectric constants. Non-limiting examples include N-alkyl pyrrolidones, such as N-ethyl pyrrolidone, diacetone alcohol, alkyl ethers, cyclic alkyl ketones, other acetyl nitriles, alcohols, and mixtures thereof. Less polar or dielectric solvents which would likely be less efficient would include non-polar hydrocarbons.

Specific examples of the methods encompassed by this invention are:

A method for reducing or removing stains from a stained fabric comprising the steps of:

- a) contacting, in the presence of water or a solvent which generates heat under microwave radiation, said stained fabric with a detergent composition comprising an effective amount of a bleaching agent;
- b) subjecting said stained fabric of Step a) to microwaves for a sufficient period to reduce or remove said stain from said fabric.

A method for deodorizing fabric having an odor, said method comprises:

- a) contacting, in the presence of water or a solvent which generates heat under microwave radiation, said fabric with a detergent composition comprising an effective amount of a bleaching agent;
- b) subjecting said fabric of Step a) to microwaves for a sufficient time to lessen the odors associated with said fabric.

A method for disinfecting an infected fabric, said method comprising:

- a) contacting, in the presence of water or a solvent which generates heat under microwave radiation, said fabric with a detergent composition comprising an effective amount of a bleaching agent;
- b) subjecting said fabric of Step a) to microwaves for a sufficient time to lessen the infecting agents associated with said fabric.

All percentages and proportions herein are by weight, and all references cited are hereby incorporated by reference, unless otherwise specifically indicated.

DETAILED DESCRIPTION OF THE INVENTION

Definitions - The present detergent compositions comprise an "effective amount" or a "stain removal-improving amount" of a particularly defined bleaching agent. An "effective amount" or "stain removal-improving amount" of a bleaching agent is any amount capable of measurably improving stain removal (especially of tea stains and carotenoid stains) from the fabric, when it is washed by the consumer. In general, this amount may vary quite widely.

By "tough food cleaning" herein is meant the ability to clean dried-on foods. Examples include dried on egg, and dried beef grease.

Microwaves - By microwaving herein is meant exposing said fabric treated with said compositions to microwave electromagnetic radiation. This is by any conventional means such as by placing the fabric in a typical microwave such as used in homes and microwaving the fabric for a sufficient time. Microwaves have an electromagnetic radiation wavelength of from about 1cm to about 1m, preferably from about 3cm to about 30cm, more preferably from about 11cm to about 13cm. See Aust. J. Chem., 1995, 48 [10], 1665-1692, Developments in Microwave-Assisted Organic Chemistry, by Strauss and Trainor.

Bleaching Agents

Suitable bleaching for use herein are listed below:

Diacyl Peroxide Bleaching Species - The composition of the present invention preferably contain diacyl peroxide of the general formula:



wherein R and R¹ can be the same or different and are hydrocarbyls, preferably no more than one is a hydrocarbyl chain of longer than ten carbon atoms, more preferably at least one has an aromatic nucleus.

Examples of suitable diacyl peroxides are selected from the group consisting dibenzoyl peroxide, dianisoyl peroxide, benzoyl gluaryl peroxide, benzoyl succinyl peroxide, di-(2-methybenzoyl) peroxide, diphthaloyl peroxide, dinaphthoyl peroxide, substituted dinaphthoyl peroxide, and mixtures thereof, more preferably dibenzoyl peroxide, dicumyl peroxide, diphthaloyl peroxides and mixtures thereof. A particularly preferred diacyl peroxide is dibenzoyl peroxide.

Hydrogen Peroxide Source - The compositions of the present invention may comprise a source of oxygen bleach, preferably a source of hydrogen peroxide with or without a selected bleach activator. The source of hydrogen peroxide is typically any common hydrogen-peroxide releasing salt, such as sodium perborate or sodium percarbonate. Hydrogen peroxide sources are illustrated in detail in Kirk Othmer Review on Bleaching and include the various forms of sodium perborate and sodium percarbonate and modified forms. An "effective amount" of a source of hydrogen peroxide is any amount capable of measurably improving stain removal (especially of tea and tomato stains) from the soiled fabric compared to a hydrogen peroxide source-free composition when the soiled fabric is washed by the consumer.

The preferred source of hydrogen peroxide used herein can be any convenient source, including hydrogen peroxide itself. For example, perborate, e.g., sodium perborate (any hydrate but preferably the mono- or tetra-hydrate), sodium carbonate peroxyhydrate or equivalent percarbonate salts, sodium pyrophosphate peroxy-hydrate, urea peroxyhydrate, or sodium peroxide can be used herein. Sodium perborate monohydrate and sodium percarbonate are particularly preferred. Mixtures of any convenient hydrogen peroxide sources can also be used.

Another source of hydrogen peroxide is enzymes. Examples include Lipoxidase, glucose oxidase, peroxidase, alcohol oxidases, and mixtures thereof.

Bleach Activators - Numerous conventional bleach activators are known. See for example activators referenced hereinabove in the background as well as U.S. Patent 4,915,854, issued April 10, 1990 to Mao et al, and U.S. Patent 4,412,934. Nonanoyloxybenzenesulfonate (NOBS) or acyl lactam activators may be used, and mixtures thereof with TAED can also be used. See also U.S. 4,634,551 for other typical conventional bleach activators. Also known are amido-derived bleach activators of the formulae: $R^1N(R^5)C(O)R^2C(O)L$ or $R^1C(O)N(R^5)R^2C(O)L$ wherein R^1 is an alkyl group containing from about 6 to about 12 carbon atoms, R^2 is an alkylene containing

from 1 to about 6 carbon atoms, R^5 is H or alkyl, aryl, or alkaryl containing from about 1 to about 10 carbon atoms, and L is any suitable leaving group. Further illustration of bleach activators of the above formulae include (6-octanamidocaproyl)oxybenzenesulfonate, (6-nonanamidocaproyl)-oxybenzenesulfonate, (6-decanamidocaproyl)oxybenzenesulfonate, and mixtures thereof as described in U.S. Patent 4,634,551. Another class of bleach activators comprises the benzoxazin-type activators disclosed by Hodge et al in U.S. Patent 4,966,723, issued October 30, 1990. Still another class of bleach activators includes acyl lactam activators such as octanoyl caprolactam, 3,5,5-trimethylhexanoyl caprolactam, nonanoyl caprolactam, decanoyl caprolactam, undecenoyl caprolactam, octanoyl valerolactam, decanoyl valerolactam, undecenoyl valerolactam, nonanoyl valerolactam, 3,5,5-trimethyl-hexanoyl valerolactam, t-butylbenzoylcaprolactam, t-butylbenzoylvalerolactam and mixtures thereof. The present compositions can optionally comprise aryl benzoates, such as phenyl benzoate, and acety triethyl citrate.

Quaternary Substituted Bleach Activators - The present compositions can also comprise quaternary substituted bleach activators (QSBA). QSBA's herein typically have the formula $E-[Z]_n-C(O)-L$, wherein group E is referred to as the "head", group Z is referred to as the "spacer" (n is 0 or 1, i.e., this group may be present or absent, though its presence is generally preferred) and L is referred to as the "leaving group". These compounds generally contain at least one quaternary substituted nitrogen moiety, which can be contained in E, Z or L. More preferably, a single quaternary nitrogen is present and it is located in group E or group Z. In general, L is a leaving group, the pKa of the corresponding carbon acid (HL) of which can lie in the general range from about 5 to about 30, more preferably, from about 10 to about 20, depending upon the hydrophilicity of the QSBA. pKa's of leaving groups are further defined in U.S. Pat. No. 4,283,301.

Preferred QSBA's herein are water-soluble but have a tendency to partition to a definite extent into surfactant micelles, especially into micelles of nonionic surfactants.

Leaving groups and solubilizing tendencies of quaternary moieties which can be present in the QSBA's are further illustrated in U.S. 4,539,130, Spt. 3, 1985 incorporated by reference. This patent also illustrates QSBA's in which the quaternary moiety is present in the leaving group L.

British Pat. 1,382,594, published Feb. 5, 1975, discloses a class of QSBA's found suitable for use herein. In these compounds, Z is a poly(methylene) or oligo(methylene) moiety, i.e., the spacer is aliphatic, and the quaternary moiety is E. U.S. 4,818,426 issued Apr. 4., 1989 discloses another class of QSBA's suitable for use herein. These compounds are quaternary ammonium carbonate esters wherein, with reference to the

above formula, the moiety Z is attached to E via a carbon atom but is attached to the carbonyl moiety through a linking oxygen atom. These compounds are thus quaternary ammonium carbonate esters. The homologous compounds wherein the linking oxygen atom is absent from Z are likewise known and are useful herein. See, for example, U.S. 5,093,022 issued March 3, 1992 and U.S. 4,904,406, issued Feb. 27, 1990. Additionally, QSBA's are described in EP 552,812 A1 published July 28, 1993, and in EP 540,090 A2, published May 5, 1993.

Chlorine Bleach - Any chlorine bleach typically known in the art is suitable for use herein. Preferred chlorine bleaches for use herein include sodium hypochlorite, lithium hypochlorite, calcium hypochlorite, chlorinated trisodium phosphates, and mixtures thereof. For more about chlorine bleaches see *Surfactant Science Series*, Vol. 5, Part II, pages 520-26.

Other Ingredients - Detergent ingredients or adjuncts optionally included in the instant compositions can include one or more materials for assisting or enhancing cleaning performance, treatment of the fabric to be cleaned, or designed to improve the aesthetics or ease of manufacture of the compositions. Other adjuncts which can also be included in compositions of the invention at their conventional art-established levels, generally from 0% to about 20% of the composition, preferably at from about 0.1% to about 10%, include one or more processing aids, water, color speckles, dyes, fillers, bleach-compatible enzymes, germicides, alkalinity sources, hydrotropes, stabilizers, perfumes, solubilizing agents, carriers. In general, materials used for the production of detergent compositions herein are preferably checked for compatibility with the essential ingredients used herein.

In the preferred embodiments, additional ingredients such as water-soluble silicates (useful to provide alkalinity and assist in controlling corrosion), dispersant polymers (which modify and inhibit crystal growth of calcium and/or magnesium salts), chelants (which control transition metals), builders such as citrate (which help control calcium and/or magnesium and may assist buffering action), and alkalis (to adjust pH) are present. Additional bleach-improving materials such as bleach catalysts may be added.

Solvent - The solvent of the present invention is of the type which the diacyl peroxide will dissolve in. The preferred solvents are selected based upon the solubility parameter value of the diacyl peroxide employed. The solubility parameter value of a compound is available from literature sources such as Polymer Handbook. Values obtained by experiments are preferred. If the solubility parameter value is not available in the literature, the value can be calculated by using any of the methods described by Robert F. Fedor's article "A Method of Estimating Both the Solubility Parameters &

Molar Volumes of Liquids", Polymer Engineering & Science, February, 1974, Vol 14, No. 2. Once the solubility parameter value is obtained of the diacyl peroxide, solvents are selected having a solubility parameter which fall within the diacyl peroxide solubility parameter.

Said solvent is preferably selected from the group consisting of N-alkyl pyrrolidones, such as N-ethyl pyrrolidone, diacetone alcohol, long chain (greater than C₆) alkyl ethers, cyclic alkyl ketones, and mixtures thereof. Amines, ethers and short chain (less than C₆) primary and secondary alcohols are preferably not present. Without being limited by theory, it is believed that the presence of these compounds may introduce stability problems. Thus, when diacyl peroxide and solvent are present in the compositions of this invention, it is further preferable that the amount of amine, ether, or primary or secondary alcohol be limited to no more than about 5%, preferably no more than about 3%, by weight of the composition.

Surfactants - Nonlimiting examples of surfactants useful herein include the conventional C₁₁-C₁₈ alkyl benzene sulfonates ("LAS") and primary, branched-chain and random C₁₀-C₂₀ alkyl sulfates ("AS"), the C₁₀-C₁₈ secondary (2,3) alkyl sulfates of the formula CH₃(CH₂)_x(CHOSO₃-M⁺)CH₃ and CH₃(CH₂)_y(CHOSO₃-M⁺)CH₂CH₃ where x and (y + 1) are integers of at least about 7, preferably at least about 9, and M is a water-solubilizing cation, especially sodium, unsaturated sulfates such as oleyl sulfate, the C₁₀-C₁₈ alkyl alkoxy sulfates ("AE_xS"; especially EO 1-7 ethoxy sulfates), C₁₀-C₁₈ alkyl alkoxy carboxylates (especially the EO 1-5 ethoxycarboxylates), the C₁₀-C₁₈ glycerol ethers, the C₁₀-C₁₈ alkyl polyglycosides and their corresponding sulfated polyglycosides, and C₁₂-C₁₈ alpha-sulfonated fatty acid esters.

If desired, the conventional nonionic and amphoteric surfactants such as the C₁₂-C₁₈ alkyl ethoxylates ("AE") including the so-called narrow peaked alkyl ethoxylates and C₆-C₁₂ alkyl phenol alkoxyates (especially ethoxylates and mixed ethoxy/propoxy), C₁₂-C₁₈ betaines and sulfobetaines ("sultaines"), and the like, can also be included in the overall compositions. The C₁₀-C₁₈ N-alkyl polyhydroxy fatty acid amides can also be used. Typical examples include the C₁₂-C₁₈ N-methylglucamides. See WO 9,206,154. Other sugar-derived surfactants include the N-alkoxy polyhydroxy fatty acid amides, such as C₁₀-C₁₈ N-(3-methoxypropyl) glucamide. The N-propyl through N-hexyl C₁₂-C₁₈ glucamides can be used for low sudsing. C₁₀-C₂₀ conventional soaps may also be used. If high sudsing is desired, the branched-chain C₁₀-C₁₆ soaps may be used. Mixtures of anionic and nonionic surfactants are especially useful. Other conventional useful surfactants are listed in standard texts.

Preferably, anionic surfactants are used herein. Without being limited by theory, it is believed that the use of anionic surfactants maximizes both cleaning performance and removal of residual bleach from the fabric being treated.

One example of a group of surfactants suitable for use herein are those selected from the group consisting of alkyl ether sulfate, long chain (greater than about C7) alkyl ethoxylate, linear alkyl benzene sulfonate (LAS), alkyl (ether) carboxylates, alkyl polyglucoside (APG), and mixtures thereof.

Thickeners - Thickeners for use herein are selected from clay, polycarboxylates, such as Polygel®, gums, carboxymethyl cellulose, polyacrylates, and mixtures thereof. The preferred clay type herein has a double-layer structure. The clay may be naturally occurring, e.g., Bentonites, or artificially made, e.g., Laponite®. Laponite® is supplied by Southern Clay Products, Inc. See *The Chemistry and Physics of Clays*, Grimshaw, 4th ed., 1971, pages 138-155, Wiley-Interscience.

Bleach catalysts - If desired, detergent compositions herein may additionally incorporate a catalyst or accelerator to further improve bleaching or starchy soil removal. Any suitable bleach catalyst can be used. The compositions will comprise from about 0.0001% to about 0.1% by weight of bleach catalyst.

Typical bleach catalysts comprise a transition-metal complex, for example one wherein the metal co-ordinating ligands are quite resistant to labilization and which does not deposit metal oxides or hydroxides to any appreciable extent under the conditions of cleaning herein. Such catalyst compounds often have features of naturally occurring compounds such as enzymes but are principally provided synthetically. Highly preferred accelerators include, for example, the cobalt 3+ catalysts, especially $\{\text{Co}(\text{NH}_3)_5\text{Cl}\}^{2+}$ or equivalents thereof with various alternate donor ligands. Such complexes include those formerly disclosed for use in laundry compositions in U.S. Pat. 4,810,410 to Diakun et al, issued March 7, 1989. The active species thereof is believed to be $\{\text{Co}(\text{NH}_3)_5(\text{OOH})\}^{2+}$ and is disclosed in J. Chem. Soc. Faraday Trans., 1994, Vol. 90, 1105-1114. Alternate catalysts or accelerators are the noncobalt transition metal complexes disclosed in this reference, especially those based on Mo(VI), Ti(IV), W(VI), V(V) and Cr(VI) although alternate oxidation states and metals may also be used. Such catalysts include manganese-based catalysts disclosed in U.S. Pat. 5,246,621, U.S. 5,244,594; U.S. 5,194,416; U.S. 5,114,606; and EP Nos. 549,271 A1, 549,272 A1, 544,440 A2, and 544,490 A1; preferred examples of these catalysts include $\text{Mn}^{\text{IV}}_2(\mu\text{-O})_3(\text{TACN})_2(\text{PF}_6)_2$, $\text{Mn}^{\text{III}}_2(\mu\text{-O})_1(\mu\text{-OAc})_2(\text{TACN})_2(\text{ClO}_4)_2$, $\text{Mn}^{\text{IV}}_4(\mu\text{-O})_6(\text{TACN})_4(\text{ClO}_4)_4$, $\text{Mn}^{\text{III}}\text{Mn}^{\text{IV}}_4-(\mu\text{-O})_1(\mu\text{-OAc})_2(\text{TACN})_2-(\text{ClO}_4)_3$, $\text{Mn}^{\text{IV}}-(\text{TACN})-(\text{OCH}_3)_3(\text{PF}_6)$, and mixtures thereof wherein TACN is trimethyl-1,4,7-

triazacyclononane or an equivalent macrocycle; though alternate metal-co-ordinating ligands as well as mononuclear complexes are also possible and monometallic as well as di- and polymetallic complexes and complexes of alternate metals such as iron or ruthenium are all within the present scope. Other metal-based bleach catalysts include those disclosed in U.S. Pat. 4,430,243 and U.S. Pat. 5,114,611. The use of manganese with various complex ligands to enhance bleaching is also reported in the following United States Patents: 4,728,455; 5,284,944; 5,246,612; 5,256,779; 5,280,117; 5,274,147; 5,153,161; and 5,227,084.

Transition metals may be precomplexed or complexed *in-situ* with suitable donor ligands selected in function of the choice of metal, its oxidation state and the denticity of the ligands. Other complexes which may be included herein are those of U.S. Application Ser. No. 08/210,186, filed March 17, 1994. Other suitable transition metals in said transition-metal-containing bleach catalysts include iron, cobalt, ruthenium, rhodium, iridium, and copper.

Builders - Detergent builders can optionally be included in the compositions herein to assist in controlling mineral hardness. Inorganic as well as organic builders can be used. Builders are typically used in fabric laundering compositions to assist in the removal of particulate soils.

The level of builder can vary widely depending upon the end use of the composition and its desired physical form. When present, the compositions will typically comprise at least about 1% builder. Liquid formulations typically comprise from about 5% to about 50%, more typically about 5% to about 30%, by weight, of detergent builder. Lower or higher levels of builder, however, are not meant to be excluded.


Inorganic or P-containing detergent builders include, but are not limited to, the alkali metal, ammonium and alkanolammonium salts of polyphosphates (exemplified by the tripolyphosphates, pyrophosphates, and glassy polymeric meta-phosphates), phosphonates, phytic acid, silicates, carbonates (including bicarbonates and sesquicarbonates), sulphates, and aluminosilicates.

Examples of silicate builders are the alkali metal silicates, particularly those having a $\text{SiO}_2:\text{Na}_2\text{O}$ ratio in the range 1.6:1 to 3.2:1 and layered silicates, such as the layered sodium silicates described in U.S. Patent 4,664,839, issued May 12, 1987 to H. P. Rieck. NaSKS-6 is the trademark for a crystalline layered silicate marketed by Hoechst (commonly abbreviated herein as "SKS-6"). NaSKS-6 can be prepared by methods such as those described in German DE-A-3,417,649 and DE-A-3,742,043. Other layered silicates, such as those having the general formula $\text{NaMSi}_x\text{O}_{2x+1} \cdot y\text{H}_2\text{O}$ wherein M is

sodium or hydrogen, x is a number from 1.9 to 4, preferably 2, and y is a number from 0 to 20 can be used herein. Various other layered silicates from Hoechst include NaSKS-5, NaSKS-7 and NaSKS-11, as the alpha, beta and gamma forms.


Examples of carbonate builders are the alkaline earth and alkali metal carbonates as disclosed in German Patent Application No. 2,321,001 published on November 15, 1973.

Aluminosilicate builders may be useful in the present invention. Aluminosilicate builders include those having the empirical formula:



wherein z and y are integers of at least 6, the molar ratio of z to y is in the range from 1.0 to about 0.5, and x is an integer from about 15 to about 264.

Useful aluminosilicate ion exchange materials are commercially available. A method for producing aluminosilicate ion exchange materials is disclosed in U.S. Patent 3,985,669, Krummel, et al, issued October 12, 1976. Preferred synthetic crystalline aluminosilicate ion exchange materials useful herein are available under the designations Zeolite A, Zeolite P (B), Zeolite MAP and Zeolite X. In an especially preferred embodiment, the crystalline aluminosilicate ion exchange material has the formula:



wherein x is from about 20 to about 30, especially about 27. This material is known as Zeolite A. Dehydrated zeolites (x = 0 - 10) may also be used herein. Preferably, the aluminosilicate has a particle size of about 0.1-10 microns in diameter.

Organic detergent builders suitable for the purposes of the present invention include, but are not restricted to, a wide variety of polycarboxylate compounds. As used herein, "polycarboxylate" refers to compounds having a plurality of carboxylate groups, preferably at least 3 carboxylates. Polycarboxylate builder can generally be added to the composition in acid form, but can also be added in the form of a neutralized salt. When utilized in salt form, alkali metals, such as sodium, potassium, and lithium, or alkanolammonium salts are preferred.

Included among the polycarboxylate builders are a variety of categories of useful materials. One important category of polycarboxylate builders encompasses the ether polycarboxylates, including oxydisuccinate, as disclosed in Berg, U.S. Patent 3,128,287, issued April 7, 1964, and Lamberti et al, U.S. Patent 3,635,830, issued January 18, 1972. See also "TMS/TDS" builders of U.S. Patent 4,663,071, issued to Bush et al, on May 5, 1987. Suitable ether polycarboxylates also include cyclic compounds, particularly alicyclic compounds, such as those described in U.S. Patents 3,923,679; 3,835,163; 4,158,635; 4,120,874 and 4,102,903.

Other useful detergency builders include the ether hydroxypolycarboxylates, copolymers of maleic anhydride with ethylene or vinyl methyl ether, 1, 3, 5-trihydroxy benzene-2, 4, 6-trisulphonic acid, and carboxymethyloxysuccinic acid, the various alkali metal, ammonium and substituted ammonium salts of polyacetic acids such as ethylenediamine tetraacetic acid and nitrilotriacetic acid, as well as polycarboxylates such as mellitic acid, succinic acid, oxydisuccinic acid, polymaleic acid, benzene 1,3,5-tricarboxylic acid, carboxymethyloxysuccinic acid, and soluble salts thereof.

Citrate builders, e.g., citric acid and soluble salts thereof (particularly sodium salt), are polycarboxylate builders of importance for liquid detergent formulations due to their availability from renewable resources and their biodegradability. Oxydisuccinates are also especially useful in such compositions and combinations.

Also suitable in the compositions of the present invention are the 3,3-dicarboxy-4-oxa-1,6-hexanedioates and the related compounds disclosed in U.S. Patent 4,566,984, Bush, issued January 28, 1986. Laurylsuccinates are the preferred builders of this group, and are described in European Patent Application 86200690.5/0,200,263, published November 5, 1986.

Other suitable polycarboxylates are disclosed in U.S. Patent 4,144,226, Crutchfield et al, issued March 13, 1979 and in U.S. Patent 3,308,067, Diehl, issued March 7, 1967. See also Diehl U.S. Patent 3,723,322.

Fatty acids, e.g., C₁₂-C₁₈ monocarboxylic acids, can also be incorporated into the compositions alone, or in combination with the aforesaid builders, especially citrate and/or the succinate builders, to provide additional builder activity.

Various alkali metal phosphates such as the well-known sodium tripolyphosphates, sodium pyrophosphate and sodium orthophosphate can be used. Phosphonate builders such as ethane-1-hydroxy-1,1-diphosphonate and other known phosphonates (see, for example, U.S. Patents 3,159,581; 3,213,030; 3,422,021; 3,400,148 and 3,422,137) can also be used.

Enzymes - Suitable enzymes include proteases, amylases, lipases, cellulases, peroxidases, and mixtures thereof of any suitable origin, such as vegetable, animal, bacterial, fungal and yeast origin. Preferred selections are influenced by factors such as pH-activity and/or stability optima, thermostability, and stability to active bleach, detergents, builders and the like. In this respect bacterial or fungal enzymes are preferred, such as bacterial amylases and proteases, and fungal cellulases.

Enzymes are normally incorporated into detergent or detergent additive compositions at levels sufficient to provide a "cleaning-effective amount". The term "cleaning effective amount" refers to any amount capable of producing a cleaning, stain

removal, soil removal, whitening, deodorizing, or freshness improving effect on fabric. In practical terms for current commercial preparations, the compositions herein may comprise from 0.001% to 5%, preferably 0.01%-1% by weight of a commercial enzyme preparation. Protease enzymes are usually present in such commercial preparations at levels sufficient to provide from 0.005 to 0.1 Anson units (AU) of activity per gram of composition.

The preparation of protease enzyme and analogous enzymes is described in GB 1,243,784 to Novo. Other suitable proteases include ALCALASE® and SAVINASE® from Novo and MAXATASE® from International Bio-Synthetics, Inc., The Netherlands; as well as Protease A as disclosed in EP 130,756 A, January 9, 1985 and Protease B as disclosed in EP 303,761 A, April 28, 1987 and EP 130,756 A, January 9, 1985. See also a high pH protease from *Bacillus* sp. NCIMB 40338 described in WO 9318140 A to Novo. Enzymatic detergents comprising protease, one or more other enzymes, and a reversible protease inhibitor are described in WO 9203529 A to Novo. Other preferred proteases include those of WO 9510591 A to Procter & Gamble. When desired, a protease having decreased adsorption and increased hydrolysis is available as described in WO 9507791 to Procter & Gamble. A recombinant trypsin-like protease for detergents suitable herein is described in WO 9425583 to Novo.

Amylases suitable herein, especially for, but not limited to automatic dishwashing purposes, include, for example, α -amylases described in GB 1,296,839 to Novo; RAPIDASE®, International Bio-Synthetics, Inc. and TERMAMYL®, Novo. FUNGAMYL® from Novo is especially useful. Engineering of enzymes for improved stability, e.g., oxidative stability, is known. See, for example *J. Biological Chem.*, Vol. 260, No. 11, June 1985, pp 6518-6521. Preferred amylases include (a) an amylase according to the hereinbefore incorporated WO 9402597, Novo, Feb. 3, 1994. Other amylases include variants having additional modification in the immediate parent as described in WO 9510603 A and are available from the assignee, Novo, as DURAMYL®. Other particularly preferred oxidative stability enhanced amylase include those described in WO 9418314 to Genencor International and WO 9402597 to Novo.

Cellulases usable herein include those disclosed in U.S. 4,435,307, Barbesgaard et al, March 6, 1984. Suitable cellulases are also disclosed in GB-A-2.075.028; GB-A-2.095.275 and DE-OS-2.247.832. CAREZYME® (Novo) is especially useful. See also WO 9117243 to Novo.

Suitable lipase enzymes for detergent usage include those produced by microorganisms of the *Pseudomonas* group, such as *Pseudomonas stutzeri* ATCC 19.154, as disclosed in GB 1,372,034. See also lipases in Japanese Patent Application

53,20487, laid open Feb. 24, 1978. Other suitable commercial lipases include Amano-CES, lipases ex *Chromobacter viscosum*, e.g. *Chromobacter viscosum* var. *lipolyticum* NRRLB 3673 from Toyo Jozo Co., Tagata, Japan; *Chromobacter viscosum* lipases from U.S. Biochemical Corp., U.S.A. and Disoynth Co., The Netherlands, and lipases ex *Pseudomonas gladioli*. LIPOLASE® enzyme derived from *Humicola lanuginosa* and commercially available from Novo, see also EP 341,947, is a preferred lipase for use herein. Lipase and amylase variants stabilized against peroxidase enzymes are described in WO 9414951 A to Novo. See also WO 9205249 and RD 94359044.

Cutinase enzymes suitable for use herein are described in WO 8809367 A to Genencor.

Peroxidase enzymes may be used in combination with oxygen sources, e.g., percarbonate, perborate, hydrogen peroxide, etc., for "solution bleaching" or prevention of transfer of dyes or pigments removed from fabrics during the wash to other substrates present in the wash solution. Known peroxidases include horseradish peroxidase, ligninase, and haloperoxidases such as chloro- or bromo-peroxidase. Peroxidase-containing detergent compositions are disclosed in WO 89099813 A, October 19, 1989 to Novo and WO 8909813 A to Novo.

A range of enzyme materials and means for their incorporation into synthetic detergent compositions is also disclosed in WO 9307263 A and WO 9307260 A to Genencor International, WO 8908694 A to Novo, and U.S. 3,553,139, January 5, 1971 to McCarty et al. Enzymes are further disclosed in U.S. 4,101,457, Place et al, July 18, 1978, and in U.S. 4,507,219, Hughes, March 26, 1985. Enzyme materials useful for liquid detergent formulations, and their incorporation into such formulations, are disclosed in U.S. 4,261,868, Hora et al, April 14, 1981. Enzymes for use in detergents can be stabilised by various techniques. Enzyme stabilisation techniques are disclosed and exemplified in U.S. 3,600,319, August 17, 1971, Gedge et al, EP 199,405 and EP 200,586, October 29, 1986, Venegas. Enzyme stabilisation systems are also described, for example, in U.S. 3,519,570. A useful *Bacillus*, sp. AC13 giving proteases, xylanases and cellulases, is described in WO 9401532 A to Novo.

Enzyme Stabilizing System - Enzyme-containing, including but not limited to, liquid compositions, herein may comprise from about 0.001% to about 10%, preferably from about 0.005% to about 8%, most preferably from about 0.01% to about 6%, by weight of an enzyme stabilizing system. Such stabilizing systems can, for example, comprise calcium ion, boric acid, propylene glycol, short chain carboxylic acids, boronic acids, and mixtures thereof, and are designed to address different stabilization problems

depending on the type and physical form of the detergent composition. See Severson, U.S. 4,537,706 for a review of Borate stabilizers.

Stabilizing systems may further comprise from 0 to about 10%, preferably from about 0.01% to about 6% by weight, of chlorine bleach scavengers, added to prevent chlorine bleach species present in many water supplies from attacking and inactivating the enzymes, especially under alkaline conditions. Suitable chlorine scavenger anions are widely known and readily available, and, if used, can be salts containing ammonium cations with sulfite, bisulfite, thiosulfite, thiosulfate, iodide, etc. Antioxidants such as carbamate, ascorbate, etc., organic amines such as ethylenediaminetetracetic acid (EDTA) or alkali metal salt thereof, monoethanolamine (MEA), and mixtures thereof can likewise be used. Other conventional scavengers such as bisulfate, nitrate, chloride, sources of hydrogen peroxide such as sodium perborate tetrahydrate, sodium perborate monohydrate and sodium percarbonate, as well as phosphate, condensed phosphate, acetate, benzoate, citrate, formate, lactate, malate, tartrate, salicylate, etc., and mixtures thereof can be used if desired.

Material Care Agents - The present compositions may optionally contain as corrosion inhibitors and/or anti-tarnish aids one or more material care agents such as silicates. Material Care Agents are preferred especially in countries where electroplated nickel silver and sterling silver are common in domestic flatware, or when aluminium protection is a concern and the composition is low in silicate. Material care agents include bismuth salts, transition metal salts such as those of manganese, certain types of paraffin, triazoles, pyrazoles, thiols, mercaptans, aluminium fatty acid salts, and mixtures thereof and are preferably incorporated at low levels, e.g., from about 0.01% to about 5% of the composition. A preferred paraffin oil is a predominantly branched aliphatic hydrocarbon comprising from about 20 to about 50, more preferably from about 25 to about 45, carbon atoms with a ratio of cyclic to noncyclic hydrocarbons of about 32 to 68 sold by Wintershall, Salzbergen, Germany as WINOG 70®. $\text{Bi}(\text{NO}_3)_3$ may be added. Other corrosion inhibitors are illustrated by benzotriazole, thiols including thionaphthol and thioanthranol, and finely divided aluminium fatty acid salts. All such materials will generally be used judiciously so as to avoid producing spots or films on glassware or compromising the bleaching action of the compositions. For this reason, it may be preferred to formulate without mercaptan anti-tarnishes which are quite strongly bleach-reactive or common fatty carboxylic acids which precipitate with calcium.

Chelating Agents - The detergent compositions herein may also optionally contain one or more iron and/or manganese chelating agents. Such chelating agents can be selected from the group consisting of amino carboxylates, amino phosphonates,

polyfunctionally-substituted aromatic chelating agents and mixtures therein, all as hereinafter defined.

Amino carboxylates useful as optional chelating agents include ethylenediaminetetracetates, N-hydroxyethylethylenediaminetriacetates, nitrilotriacetates, ethylenediamine tetrapropionates, triethylenetetra-amine-hexacetates, diethylenetriaminepentaacetates, and ethanoldiglycines, alkali metal, ammonium, and substituted ammonium salts therein and mixtures therein.

Polyfunctionally-substituted aromatic chelating agents are also useful in the compositions herein. See U.S. Patent 3,812,044, issued May 21, 1974, to Connor et al. Preferred compounds of this type in acid form are dihydroxydisulfobenzenes such as 1,2-dihydroxy-3,5-disulfobenzene. A preferred biodegradable chelator for use herein is ethylenediamine disuccinate ("EDDS"), especially the [S,S] isomer as described in U.S. Patent 4,704,233, November 3, 1987, to Hartman and Perkins.

If utilized, these chelating agents will generally comprise from about 0.1% to about 10% by weight of the detergent compositions herein. More preferably, if utilized, the chelating agents will comprise from about 0.1% to about 3.0% by weight of such compositions.

Polymeric Dispersing Agents - Polymeric dispersing agents can advantageously be utilized at levels from about 0.1% to about 7%, by weight, in the compositions herein, especially in the presence of zeolite and/or layered silicate builders. Suitable polymeric dispersing agents include polymeric polycarboxylates and polyethylene glycols, although others known in the art can also be used. It is believed, though it is not intended to be limited by theory, that polymeric dispersing agents enhance overall detergent builder performance, when used in combination with other builders (including lower molecular weight polycarboxylates) by crystal growth inhibition, particulate soil release peptization, and anti-redeposition.

Polymeric Soil Release Agent - Any polymeric soil release agent known to those skilled in the art can optionally be employed in the compositions and processes of this invention. Polymeric soil release agents are characterized by having both hydrophilic segments, to hydrophilize the surface of hydrophobic fibers, such as polyester and nylon, and hydrophobic segments, to deposit upon hydrophobic fibers and remain adhered thereto through completion of washing and rinsing cycles and, thus, serve as an anchor for the hydrophilic segments. This can enable stains occurring subsequent to treatment with the soil release agent to be more easily cleaned in later washing procedures.

If utilized, soil release agents will generally comprise from about 0.01% to about 10.0%, by weight, of the detergent compositions herein, typically from about 0.1% to about 5%, preferably from about 0.2% to about 3.0%.

Without being limited by theory, it is believed that the presence of solvents may act to dissolve the diacyl peroxide and thus reduce the abrasive character of the composition. Thus, it is preferable that the solvent be limited to no more than about 4%, preferably no more than about 2%, by weight of the composition. However, solvents may be present to assist in the formation of suitable particles for use herein.

Brightener - Any optical brighteners or other brightening or whitening agents known in the art can be incorporated at levels typically from about 0.05% to about 1.2%, by weight, into the detergent compositions herein. Commercial optical brighteners which may be useful in the present invention can be classified into subgroups, which include, but are not necessarily limited to, derivatives of stilbene, pyrazoline, coumarin, carboxylic acid, methinecyanines, dibenzothiphen-5,5-dioxide, azoles, 5- and 6-membered-ring heterocycles, and other miscellaneous agents. Examples of such brighteners are disclosed in "The Production and Application of Fluorescent Brightening Agents", M. Zahradnik, Published by John Wiley & Sons, New York (1982).

Methods for producing diacyl peroxide particles for use in the compositions herein wherein an abrasive particle is desired may include any particle making process commonly known in the art, including shear mixing. The diacyl particles for use herein can range in size from sub-micron (0.1) to about 100 microns. A preferred range is from about 1 to about 20 microns. Another process for making particles follows:

Process Description - The diacyl peroxide raw material particles are dissolved in an appropriate solvent (n-ethylpyrrolidone) and added to the rest of the formulation (primarily water, surfactant and thickener) with stirring. This procedure results in the *in situ* precipitation of the diacyl peroxide particles, resulting in a dispersion of small homogeneous particles ranging in size of from about 1 to about 20 microns.

Procedure for preparation of in situ particles: Laponite (33g, 6% active) is dispersed in tap water (100g) with stirring. Sodium Alkylethoxy sulfate (14g, 70% active) is stirred into the Laponite dispersion and Sodium bicarbonate (1g, 100% active) is added. In a separate container, Benzoyl peroxide (2g, 75% active) is dissolved in N-ethylpyrrolidone (10g, 100% active) with stirring. This benzoyl peroxide solution is then poured into the Laponite and surfactant solution with stirring. The mixture immediately turns cloudy and results in a homogeneous dispersion of 10-50 micron benzoyl peroxide particles.

Product/Instructions - This invention also encompasses the inclusion of instructions on the use of the cleaning composition with the package containing the cleaning compositions herein or with other forms of advertising associated with the sale or use of the cleaning compositions. The instructions may be included in any manner typically used by consumer product manufacturing or supply companies. Examples include providing instructions on a label attached to the container holding the composition; on a sheet either attached to the container or accompanying it when purchased; or in advertisements, demonstrations, and/or other written or oral instructions which may be connected to the purchase or use of the cleaning compositions.

Specifically the instructions will include a description of the use of the cleaning composition in connection with microwaving. The instructions, for instance, may additionally include information relating to the length of microwaving time; the recommended settings on the microwave; the recommended amount of treating composition to apply to the substrate, if soaking or rubbing is appropriate to the substrate; the recommended amount of water, if any, to apply to the substrate before and after treatment; other recommended treatment to accompany the microwave application.

A product comprising a treating composition comprising a bleaching agent and instructions for use of the treating composition, said instructions include the steps of:

- contacting a substrate, in the presence of water or a solvent which generates heat under microwave radiation, (either in the product or supplied separately) with said treating composition; and
- subjecting said substrate to microwaves for a sufficient period to effectively treat said substrate.

EXAMPLE I

Non-aqueous Liquid Detergent Composition with Bleach	
<u>Component</u>	<u>Wt% Active</u>
LAS powder	20.36
C12-14EO+5 alcohol ethoxylate	18.8
Butoxy-Propoxy-Propanol	19
Sodium citrate dihydrate	4
Citrate Coated Nonanoyloxybenzenesulfonate	8.5
Sodium carbonate	12
Maleic-acrylic copolymer	12
DTPA	1

Sodium Perborate	3
Suds Suppressor	0.03
Perfume	0.5
Titanium Dioxide	0.5
Brightener	0.3
Total	to 100%

DTPA is diethyltriamine pentaacetic acid

LAS powder is a combination of (by weight): Na linear alkylbenzenesulfonate 85%; sulfate 11%; sulfosuccinate 2%; water 2.5%; balance to 100% of unreacted byproducts.

One-fourth cup of the above formula is applied directly to a dampened 0.3 meter by 0.3 meter fabric sample and microwaved for approximately 1 minute on high setting in a conventional, household microwave. One-fourth cup of the above formula is also applied directly to a dampened fabric sample and allowed to set for approximately one minute. Both fabric samples are washed in a conventional laundry manner at 32°C. The results are shown below.

Stain Set	No microwave	Microwaved 1 minute)
Greasy Foods (butter, burnt lard)	73.3	94.0
Hydrophobic Stains (tomato, curry, carotene)	79.8	89.1
Hydrophilic Stains (tea, wine, coffee)	51.5	83.3
Particulate Stains (clays, peat, make-up)	63.5	62.1
Enzymatic Stains (chocolate, blood, grass)	78.5	66.9

The microwaved composition provides significant performance benefits in 3 of the 5 stain categories.

EXAMPLE II

Component	A (% Wt)	B (% Wt)	C (% Wt)
Surfactants			
Alkyl Sulfate	6	3	2
EO/PO Block Polymer	--	0.3	--
Neodol 45-7	--	--	6.5
Neodol 23-3	--	--	8.5
Bleaches			
Acetyl Triethylcitrate	--	--	3.5
Benzoyl Peroxide	1.5	1.5	--
Hydrogen Peroxide	--	--	4
Solvents/Buffers			
N-Ethyl pyrrolidone	7	--	--
Triethanolamine	--	--	1.2
Sodium bicarbonate	0.8	0.8	--

Thickeners			
Laponite Clay	1.2	1.2	--
Water	Balance	Balance	Balance
pH	7	7	5

EXAMPLE III

<u>Ingredient</u>	<u>Acyl peroxide</u>	<u>Acyl peroxide</u>
C ₁₀ alkyl ethoxylate (avg. ethoxy of 10)	5	-
C ₁₂₋₁₃ alkyl ether sulfate (avg. ethoxy of 1)	14	12
Magnesium chloride hexahydrate	0.3	-
Magnesium silicate ¹	2	1.5
Potassium bicarbonate	1	1
Alkyl pyrrolidinone	15	7
Acyl peroxide ²	2	-
Perfume	0.18	-
Other (water, dye etc.)	to 100%	to 100%
Performance, Stain Removal		
Tomato on plastic	very good	very good
Tomato on cotton	very good	very good
Tea on ceramic	very good	very good

¹ Commercially available as Laponite RD®

² Acyl Peroxides selected from dibenzoyl peroxide, dianisoyl peroxide, benzoyl gluaryl peroxide, benzoyl succinyl peroxide, di-(2-methybenzoyl) peroxide, diphtaloyl peroxide, dinaphthoyl peroxide, substituted dinaphthoyl peroxide, and mixtures thereof.

Ceramic and plastic cups, bowls etc. are stained by heating tomato sauce and/or tea under consumer relevant conditions in the microwave. The stained items are washed with a conventional light duty liquid dishwashing detergent that is commercially available under typical home wash conditions. The objects remain stained by the tomato and tea. An amount of the new bleach/detergent composition is applied to the stained item until the stained item is evenly coated with the composition. The treated item is placed in a typical household microwave and microwaved on high setting for 30-45

seconds. The item is then rinsed out. The percent removal is estimated visually based on comparison with a stained control and a clean control. As can be seen in the example, in A without bleaching agent has poor stain removal ability. Examples B and C have good stain removal ability.

WHAT IS CLAIMED IS:

- 1.) A method for treating fabrics comprising the steps of:
 - a) contacting said fabrics, in the presence of water or a solvent which generates heat under microwave radiation, with a treating composition comprising an effective bleaching agent;
 - b) subjecting said fabric to microwaves for a sufficient period to effectively treat said fabric.
- 2.) A method according to Claim 1 wherein said fabric is treated for one or more of the following: stain removal, stain reducing, deodorizing, or disinfecting.
- 3.) A method according to Claim 1 wherein said bleaching agent is selected from the group consisting of:
 - a) diacyl peroxide;
 - b) a source of hydrogen peroxide and bleach activator;
 - c) a source of hydrogen peroxide;
 - c) a chlorine bleach; and
 - d) mixtures thereof.
- 4.) A method according to Claim 4 wherein said treating composition further comprises one or more of the following: surfactant, solvent, clay, water, polycarboxylate thickeners, baking soda, carbonates, phosphates, water, hydrobenzoic acid, dicarboxylic acid, siloxanes, perfumes, bleach catalysts, and mixtures thereof.
- 5.) A method according to Claim 1 wherein said sufficient time in step b) is from 1 second to 2 minutes.
- 6.) A method according to Claim 1 wherein said diacyl peroxides are selected from the group consisting dibenzoyl peroxide, dianisoyl peroxide, benzoyl gluaryl peroxide, benzoyl succinyl peroxide, di-(2-methybenzoyl) peroxide, diphthaloyl peroxide, dinaphthoyl peroxide, substituted dinaphthoyl peroxide, and mixtures thereof.

- 7.) A method according to Claim 1 wherein said treating composition is a sprayable liquid or sprayable gel detergent compositions especially effective at cleaning stains, deodorizing, and/or disinfecting fabrics, comprising by weight:

(a) from 0.1% to 60% of a bleaching agent selected from the group consisting of:

i) diacyl peroxide having the general formula:



wherein R and R1 can be the same or different;

- ii) a source of hydrogen peroxide;
 - iii) a source of hydrogen peroxide and a bleach activator;
 - iv) a chlorine bleach; and
 - v) mixtures thereof;
- (b) from 0% to 95% of a non-aqueous solvent;
- (c) from 0% to 50% of a surfactant; and
- (d) from 0% to 7% of a thickener.

- 8.) A method according to Claim 1 wherein said treating composition is a gel detergent composition especially effective at cleaning stains and tough food soil from fabrics, comprising by weight:

(a) from 0.1% to 10% of a diacyl peroxide having the general formula:



wherein R and R1 can be the same or different;

- (b) from 0% to 50% of a surfactant; and
- (c) from 0% to 7% of a thickener;

said composition having a neat pH of from 3 to 10; and such that said diacyl peroxide remains undissolved in said compositions.

- 9.) A product comprising a treating composition comprising a bleaching agent and instructions for use of the treating composition, said instructions include the steps of:

- a) contacting a fabric, in the presence of water or a solvent which generates heat under microwave radiation, with said treating composition; and
- b) subjecting said fabric to microwaves for a sufficient period to effectively treat said fabric.

THIS PAGE BLANK (USPTO)



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C11D 3/39, 3/395	A3	(11) International Publication Number: WO 98/03621 (43) International Publication Date: 29 January 1998 (29.01.98)
(21) International Application Number: PCT/US97/12215 (22) International Filing Date: 18 July 1997 (18.07.97) (30) Priority Data: 60/022,334 24 July 1996 (24.07.96) US (71) Applicant (for all designated States except US): THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): OFOSU-ASANTE, Kofi [GH/US]; 1535 Cedar Avenue, Cincinnati, OH 45224 (US). HUTTON, Howard, David [US/US]; 2745 Jessica Court, Loveland, OH 45140 (US). (74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 16 April 1998 (16.04.98)
(54) Title: METHOD FOR STAIN REMOVAL ON FABRIC WITH DETERGENT COMPOSITIONS CONTAINING BLEACH (57) Abstract <p>The present invention is in the field of liquid or gel detergents. More specifically, the invention relates to a method for removing stains and/or odors and for disinfecting fabrics by applying a liquid or gel detergent composition to the fabric and subjecting the fabric to microwaves. This method provides enhanced cleaning and improved stain removal. These cleaning compositions comprise a bleaching agent, preferably diacyl peroxide in a liquid or gel detergent formulation. These compositions in combination with the method of microwaving are also very effective in disinfecting and deodorizing fabrics.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 97/12215

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C11D3/39 C11D3/395

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C11D D06M

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DE 37 03 049 A (WFK TESTGEWEBE GMBH) 11 August 1988 see example 1 see claims 1-11 ---	1-5, 7, 9
X	EP 0 295 205 A (CIBA GEIGY AG) 14 December 1988 see claim 15 ---	1-5
A	EP 0 024 304 A (DEPPE HANS DIETER DR) 4 March 1981 see example 2 ---	1-3
A	DE 36 06 823 A (SCHUBERT WERNER) 10 September 1987 see claim 1 -----	1

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

° Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

3 February 1998

Date of mailing of the international search report

18/02/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Richards, M

INTERNATIONAL SEARCH REPORT

...for information on patent family members

International Application No

PCT/US 97/12215

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 3703049 A	11-08-88	NONE	
EP 0295205 A	14-12-88	FR 2616164 A JP 63309685 A US 4861342 A	09-12-88 16-12-88 29-08-89
EP 0024304 A	04-03-81	DE 2933498 A AT 5762 T	09-04-81 15-01-84
DE 3606823 A	10-09-87	DE 3606822 A	05-11-87